

**Table 1:** Morphometric measurements (in millimetres) of *Megaderma lyra* and *Megaderma spasma* specimens caught from the temple roost in Lamheti village in Jabalpur Madhya Pradesh

Name of the body part	<i>Megaderma lyra</i>		<i>Megaderma spasma</i>
	Male (n=1)	Male (n=1)	Female (n=1)
Head and Body length	80.2	65.2	66.3
Hind foot	15.9	14.9	14.8
Forearm	64.3	55.7	55.5
Ear	36.7	34.2	34.6
Wing span	410.4	308	311
Nose leaf	9.4	6.1	6.2

inside a temple in Karnataka, the co-existence of *M. lyra* and *M. spasma* has been rarely seen in India.

#### ACKNOWLEDGEMENTS

I extend my sincere thanks to my colleague Mr. Anant Khot of the Bombay Natural History Society (BNHS) for assisting me in field work and Mr. Mayank and Mr. Rahul of Lamheti village for guiding me to the above-mentioned bat roost. I am also grateful to Dr. Asad R. Rahmani, Director, BNHS, for his academic support to carry out this work.

#### REFERENCES

- BATES, P.J.J. & D.L. HARRISON (1997): Bats of the Indian Subcontinent. Harrison Zoological Museum, Kent, England. Pp. 51-55.  
 WROUGHTON, R.C. (1913): Report No. 6: Kanara (29-44). Bombay Natural History Society's Mammal Survey of India. *J. Bombay Nat. Hist. Soc.* 22(1): 29-66.

## 2. SECOND RECORD OF ALBINO FIVESTRIPED PALM SQUIRREL *FUNAMBULUS PENNANTI* WROUGHTON FROM UDAIPUR, RAJASTHAN<sup>1</sup>

SATIYA PRAKASH MEHRA<sup>2</sup>, JITENDRA SINGH KHARWAR<sup>3</sup> AND NARAYAN SINGH KHARWAR<sup>3,4</sup>

<sup>1</sup>Accepted September 09, 2006

<sup>2</sup>Kesar Bhawan, 16/747, Plot No. 90, B/D Saraswati Hosp., Ganeshnagar, Pahada, Udaipur 313 001, Rajasthan, India.

Email: spmehra@yahoo.com

<sup>3</sup>74, Navratan Complex, Bedla Road, Fatehpura, Udaipur 313 001, Rajasthan, India.

<sup>4</sup>Email: narayansingh78@yahoo.co.in

The Fivestriped Palm Squirrel *Funambulus pennanti* has distinct five pale strips on its grayish-brown or olive-brown body (Menon 2003; Prater 2005). A number of local races are recognized on the basis of differences in the lightness or darkness of the coat, or variations in the tones of the dorsal strips (Prater 2005). Unlike the characters mentioned in the guide books, we sighted a solitary, white Fivestriped Palm Squirrel in the residential area of north Udaipur (Rajasthan) from mid-July 2005. On close observation of the animal, we found that the Squirrel was albino with only a small brownish patch in the middle of the dorsal side of the body (Eds: photographic evidence provided); the eyes were red. This is the second case of albinism in the Fivestriped Palm Squirrel from Rajasthan. Although, the first record of albino Fivestriped from Rajasthan was by Sharma (2004); albinism in the Fivestriped Squirrel has been reported only thrice in India (Mahabal *et al.* 2005).

According to Mahabal *et al.* (2005), albinism is said to

be rare in rodents. This is also evident from the earlier works (Rajgopalan 1967; Pradhan 1975; Chaturvedi and Ghosh 1984).

The reported albino squirrel was sighted throughout a cloudy day but its movement was restricted to morning and evening hours on sunny days. This albino individual lived in a house under construction

The solitary individual was very alert. As soon as we approached, it would escape. A distance of more than 6 m was maintained while taking observations and photographing. How long it may be sighted after the house is occupied cannot be predicted.

#### ACKNOWLEDGEMENTS

We thank Dr. Satish Kumar Sharma and Dr. Pratap Singh for valuable discussions. Also, we are grateful to Mr. Praveen Sharma and Mr. Shambhu Sharma for technical assistance in photography and videography.

#### REFERENCES

- CHATURVEDI, Y. & A.K. GHOSH (1984): A case of albinism in the Fivestriped Palm Squirrel, *Funambulus pennanti* Wroughton. *Bull. Zool. Surv. India* 6(1-3): 321-322.  
 MAHABAL, A., R.M. SHARMA & M.S. PRADHAN (2005): A case of total albinism in the Fivestriped Palm Squirrel *Funambulus pennanti* Wroughton in Sindhudurg District, Maharashtra State. *J. Bombay Nat. Hist. Soc.* 102(1): 98-99.  
 MENON, V. (2003): A Field Guide to Indian Mammals. Penguin Book India (P) Ltd and Dorling Kindersley (India) Pvt. Limited, Delhi. Pp. 128.  
 PRADHAN, M.S. (1975): Studies on Bombay Rats. Ph.D. Thesis, Bombay University, Bombay. Pp. 335

PRATER, S.H. (2005): The Book of Indian Animals. 12<sup>th</sup> Reprint. Oxford University Press and Bombay Natural History Society, Mumbai. Pp. 200-201.  
SHARMA, S.K. (2004): Occurrence of Albino Common Palm Civet and

Northern Palm Squirrel in Southern Rajasthan. *Zoos' Print Journal* 19(5): 1483.

RAJGOPALAN, P.K. (1967): A case of albinism in *Rattus blanfordi* (Thomas) 1881, (Rodentia: Muridae). *J. Bombay Nat. Hist. Soc.* 64(3): 554.

### 3. OCCURRENCE OF ASIATIC BRUSHTAILED PORCUPINE *ATHERURUS MACROURUS* (LINN. 1758) AT MIZORAM, INDIA<sup>1</sup>

NARESH CHATURVEDI<sup>2</sup>, M. SAWMLIANA<sup>3</sup> AND VINOD PATIL<sup>2,4</sup>

<sup>1</sup>Accepted March 29, 2006

<sup>2</sup>Bombay Natural History Society, Hornbill House, Shaheed Bhagat Singh Road, Mumbai 400 001, Maharashtra, India.

<sup>3</sup>Chanmari (West), Aizawl, Mizoram 796 007, India.

<sup>4</sup>Email: vinodpatil72@rediffmail.com

The Asiatic Brushtailed Porcupine *Atherurus macrourus* is known to exist in Assam, India. Ellerman and Morrison-Scott (1951) had recorded its distribution as Assam (India), Tenasserim (Myanmar), China, Indo-China, Thailand and Malaysia.

According to Choudhary (1997) *Atherurus macrourus assamensis*, a subspecies of the Asiatic Brushtailed Porcupine is found in India. This is also endorsed by Molur *et al.* (1998). In the CHECKLIST OF INDIAN MAMMALS, Nameer (2000) has also mentioned Assam as its known distribution limit. Thomas (1921) had described the specimen of Asiatic Brushtailed Porcupine collected from Assam by Wells, during the Mammal Survey of India. In his report, "Scientific results from the Mammal Survey", he had described it as a new species *Atherurus assamensis*. But subsequently, it was given the status of subspecies *assamensis*. Currently, the subspecies *assamensis* is also considered as a synonym of *Atherurus*

*macrourus* (Wilson and Reeder 1993). Until now, the known distribution of the species was only Assam. Recently, one of us (M. Swamlina) sent a photograph of an animal taken at Hmuifang, Aizawl, Mizoram for identification to the BNHS; where it was identified as *Atherurus macrourus*. The occurrence of *A. macrourus* in the Aizawl district of Mizoram state is a new distribution record for the species which was so far known only from Assam. Interestingly, Mizoram and Myanmar share a common boundary.

The Bombay Natural History Society's collections has a male specimen (Reg. No. 8997) of this species collected at Tenasserim, Myanmar on 20.xii.1913.

According to Blanford (1891), this species was restricted to the east of Bay of Bengal. He further states, "the genus must have existed in the Indian Peninsula, for its teeth have been found in the Pleistocene cave-deposits of Kurnool".

#### REFERENCES

BLANFORD, W.T. (1891): The Fauna of British India, including Ceylon and Burma, Mammalia. Taylor and Francis, Red Lion Court Fleet Street, London. 617 pp.

CHOUDHARY, ANWARUDDIN (1997): Checklist of the Mammals of Assam. Revised Edition (1997), Gibbons Books with Assam Science Technology & Environment Council, 1997, 103 pp.

ELLERMAN, J.R. & T.C.S. MORRISON-SCOTT (1951): Checklist of Palaearctic and Indian Mammals, 1758-1946. British Museum of Natural History, London, 810 pp.

MOLUR S., P.O. NAMEER & S. WALKER (Eds) (1998): Report of the

Workshop "Conservation Assessment and Management Plan for Mammals of India", Zoo Outreach Organisation, Conservation Breeding Specialist Group, Coimbatore, India. 176 pp.

NAMEER, P.O. (2000): Checklist of Indian Mammals. Kerala Forest Department. 90+xxv pp.

THOMAS, O. (1921): Scientific results from Mammal Survey XXV. *J. Bombay Nat. Hist. Soc.* 27(3): 596-598.

WILSON, D.E. & D.M. REEDER (1993): Mammal Species of the World: A Taxonomic and Geographic Reference, 2nd edn., Smithsonian Institution Press, Washington, 1206 pp.

### 4. PREDATION ON CHITAL *AXIS AXIS* BY WILD PIG *SUS SCROFA* IN BANDHAVGARH NATIONAL PARK<sup>1</sup>

SATYARANJAN BEHERA<sup>2,3</sup> AND RAJENDRA PRASAD GUPTA<sup>2,4</sup>

<sup>1</sup>Accepted October 23, 2007

<sup>2</sup>Wildlife Institute of India, Chandrabani, P.O. Box 18, Dehradun 248 001, Uttarakhand, India.

<sup>3</sup>Email: sataya240437@yahoo.co.in

<sup>4</sup>Email: pr.rajn@gmail.com

On April 12, 2006, while we were at Bandhavgarh National Park to estimate Tiger population, we observed a

group of six to seven Chital *Axis axis* grazing near a water hole at around 1700 hrs. A group of 12-13 Wild Pigs